arXiv:0908.3651v2 [nlin.SI] 2 Mar 2010

Permutability of Backlund Transformations for $N = 2$ Supersymmetric Sine-Gordon

J.F. Gomes, L.H. Ymai and A.H. Zimerman

Instituto de Física Teórica, IFT-UNESP
Universidade Estadual Paulista
Caixa Postal 70532-2
01156-970 São Paulo, SP, Brazil

ABSTRACT

The permutability of two Backlund transformations is employed to construct a non linear superposition formula and to generate a class of solutions for the $N = 2$ super sine-Gordon model. We present explicitly the one and two soliton solutions.

1 Introduction

Backlund transformations reduce the order of the non-linear differential equations making the system sometimes effectively more tractable. Starting with a simple input solution, we may be able to solve for a more complicated one. In many cases, this may be very difficult to accomplish. A convenient and powerful way is to use the *permutability theorem* which provides a closed algebraic non-linear superposition formula for the solutions.

The Backlund transformation and the Permutability theorem are employed to derive a series of consistency conditions which are satisfied by soliton solutions of certain class of integrable models. Within such class, we encounter the sine-Gordon [1] and KdV [2] equations. This framework was also applied to the $N = 1$ super KdV [3] and super sinh-Gordon [4] in order to derive its soliton solutions.

The $N = 2$ super sine-Gordon model was proposed in [5] and later in [6] its algebraic structure was uncovered. Certain solutions of this model have already been constructed [7], however they were such that involve a single Grassmann parameter. In this paper we extend the non-linear superposition formulae for soliton solutions of the $N = 2$ super sine-Gordon model. These formulae are derived from the Backlund transformation proposed in [8] and the permutability condition which implies that the order of two successive Backlund transformations is irrelevant. As examples, we present explicitly the 1-and 2-soliton solutions with distinct Grassmann parameters.

Recently the Pohlmeyer reduction of AdS_nxS_n superstring models have been considered [9] which in the simple case of $n = 2$ was shown [10] to be equivalent to the $N = 2$ supersymmetric sine-Gordon.

This paper is organized as follows. In Section 2 we discuss the $N = 2$ super sine-Gordon and its Backlund Transformation. In Section 3 we apply the permutability condition to derive a closed algebraic non-linear superposition formulae involving solutions of the model. Finally in Section 4 and 5 we present the 1- and 2-soliton solutions respectively. In the appendix A we present the Backlund transformation in components. In appendices B and C we give details for the derivation of the superposition formulae.

2 $N = 2$ super sine-Gordon - Backlund Transformation

Let us start by introducing the $N = 2$ superfields [5]

$$\phi^{\pm} \;=\; \varphi^{\pm}(z^{\pm},\bar{z}^{\pm}) + \theta^{\pm}\psi^{\mp}(z^{\pm},\bar{z}^{\pm}) + \bar{\theta}^{\pm}\bar{\psi}^{\mp}(z^{\pm},\bar{z}^{\pm}) + \theta^{\pm}\bar{\theta}^{\pm}F^{\pm}(z^{\pm},\bar{z}^{\pm}),$$

where

$$z^{\pm} = z \pm \frac{1}{2}\theta^{+}\theta^{-}, \qquad \bar{z}^{\pm} \;=\; \bar{z} \pm \frac{1}{2}\bar{\theta}^{+}\bar{\theta}^{-}.$$

The superfield components $\phi^{\pm}$ can be expanded in Grassmann variables $\theta^{\pm}$ and $\bar{\theta}^{\pm}$. For instance, the component $\varphi^{\pm}(z^{\pm},\bar{z}^{\pm})$ gives rise to

$$\varphi^{\pm}(z^{\pm},\bar{z}^{\pm}) \;=\; \varphi^{\pm} \pm \frac{1}{2}\theta^{+}\theta^{-}\partial_{z}\varphi^{\pm} \pm \frac{1}{2}\bar{\theta}^{+}\bar{\theta}^{-}\partial_{\bar{z}}\varphi^{\pm} + \frac{1}{4}\theta^{+}\theta^{-}\bar{\theta}^{+}\bar{\theta}^{-}\partial_{z}\partial_{\bar{z}}\varphi^{\pm}.$$

By expanding all components of $\phi^{\pm}$,we obtain

$$\begin{aligned}\phi^{\pm} &= \varphi^{\pm}+\theta^{\pm}\psi^{\mp}+\bar{\theta}^{\pm}\bar{\psi}^{\mp}\pm\frac{1}{2}\theta^{+}\theta^{-}\partial_{z}\varphi^{\pm}\pm\frac{1}{2}\bar{\theta}^{+}\bar{\theta}^{-}\partial_{\bar{z}}\varphi^{\pm}+\theta^{\pm}\bar{\theta}^{\pm}F^{\pm}\\ &\pm\theta^{\pm}\bar{\theta}^{+}\bar{\theta}^{-}\frac{1}{2}\partial_{\bar{z}}\psi^{\mp}\pm\bar{\theta}^{\pm}\theta^{+}\theta^{-}\frac{1}{2}\partial_{z}\bar{\psi}^{\mp}+\frac{1}{4}\theta^{+}\theta^{-}\bar{\theta}^{+}\bar{\theta}^{-}\partial_{z}\partial_{\bar{z}}\varphi^{\pm}.\end{aligned}$$

We next introduce the super derivatives

$$D_{\pm}=\frac{\partial}{\partial\theta^{\pm}}+\frac{1}{2}\theta^{\mp}\partial_{z},\qquad \bar{D}_{\pm}=\frac{\partial}{\partial\bar{\theta}^{\pm}}+\frac{1}{2}\bar{\theta}^{\mp}\partial_{\bar{z}},$$

satisfying the following conditions

$$\begin{aligned}&D_{\pm}^{2}=0, &&\bar{D}_{\pm}^{2}=0,\\ &\{\bar{D}_{\pm},D_{\pm}\}=0, &&\{\bar{D}_{\pm},D_{\mp}\}=0,\\ &\{D_{+},D_{-}\}=\partial_{z}, &&\{\bar{D}_{+},\bar{D}_{-}\}=\partial_{\bar{z}}.\end{aligned}$$

The equations of motion for the supersymmetric sine-Gordon model with $N=2$ are given by [5]

$$\bar{D}_{\pm}D_{\pm}\phi^{\pm}=g\,\sin\left(\beta\phi^{\mp}\right), \tag{1}$$

where g is a mass parameter and β is the coupling constant. From now on we assume $\beta=1$ which may be re-inserted by a convenient field reparametrization. In components, the equations of motion for the $N=2$ super sine-Gordon reads,

$$\begin{aligned}F^{\pm} &= g\,\sin\varphi^{\mp},\\ \partial_{\bar{z}}\psi^{\mp} &= g\,\cos\varphi^{\mp}\bar{\psi}^{\pm},\\ \partial_{z}\bar{\psi}^{\mp} &= -g\,\cos\varphi^{\mp}\psi^{\pm},\\ \partial_{z}\partial_{\bar{z}}\varphi^{\pm} &= -g\,\cos\varphi^{\mp}F^{\mp}-g\,\sin\varphi^{\mp}\psi^{\pm}\bar{\psi}^{\pm}.\end{aligned}$$

Moreover the the chiral, ϕ^{+} and the anti-chiral, ϕ^{-} superfields satisfy the conditions

$$\bar{D}_{\pm}\phi^{\mp}=D_{\pm}\phi^{\mp}=0. \tag{2}$$

Let us now recall the Backlund transformation for the $N=2$ super sine-Gordon model [8]. For this purpose, consider the pair of first order differential equations

$$D_{+}\phi_{1}^{+} = D_{+}\phi_{2}^{+}-\frac{8}{\kappa}\mathcal{F}\cos\left(\frac{\phi_{1}^{-}+\phi_{2}^{-}}{2}\right), \tag{3}$$

$$\bar{D}_{+}\phi_{1}^{+} = -\bar{D}_{+}\phi_{2}^{+}+\kappa\mathcal{G}\cos\left(\frac{\phi_{1}^{-}-\phi_{2}^{-}}{2}\right), \tag{4}$$

where $\mathcal{F}$ and $\mathcal{G}$ are fermionic auxiliary superfields and κ is an arbitrary constant. The above equation and the condition

$$(\bar{D}_{+}D_{+}+D_{+}\bar{D}_{+})\phi_{1}^{+}=0,$$

leads to the equations of motion

$$\bar{D}_+ D_+ \phi_2^+ = g\, \sin \phi_2^-,$$

provided the superfields $\mathcal{F}$ and $\mathcal{G}$ satisfy

$$\bar{D}_+ \mathcal{F} = -\frac{\kappa g}{4} \sin\left(\frac{\phi_1^- - \phi_2^-}{2}\right), \qquad D_+ \mathcal{G} = -\frac{2g}{\kappa} \sin\left(\frac{\phi_1^- + \phi_2^-}{2}\right). \tag{5}$$

In a similar way,

$$\begin{aligned} D_- \phi_1^- &= D_- \phi_2^- + \lambda \mathcal{G} \cos\left(\frac{\phi_1^+ + \phi_2^+}{2}\right), & (6) \\ \bar{D}_- \phi_1^- &= -\bar{D}_- \phi_2^- - \frac{8}{\lambda} \mathcal{F} \cos\left(\frac{\phi_1^+ - \phi_2^+}{2}\right), & (7) \end{aligned}$$

where λ is another arbitrary constant. Together with the condition

$$(\bar{D}_- D_- + D_- \bar{D}_-) \phi_1^- = 0,$$

yields

$$\bar{D}_- D_- \phi_2^- = g\, \sin \phi_2^+,$$

provided $\mathcal{G}$ and $\mathcal{F}$ satisfy

$$\bar{D}_- \mathcal{G} = \frac{2g}{\lambda} \sin\left(\frac{\phi_1^+ - \phi_2^+}{2}\right), \qquad D_- \mathcal{F} = \frac{\lambda g}{4} \sin\left(\frac{\phi_1^+ + \phi_2^+}{2}\right). \tag{8}$$

Acting with D_+ in eqn. (3), $\bar{D}_+$ in (4), D_- in (6) and $\bar{D}_-$ in (7) we find

$$D_+ \mathcal{F} = 0, \qquad \bar{D}_+ \mathcal{G} = 0, \qquad D_- \mathcal{G} = 0, \qquad \bar{D}_- \mathcal{F} = 0. \tag{9}$$

These last conditions allows us to rewrite the fermionic superfields into two distinct manners, i.e.,

$$\begin{aligned} \mathcal{F} &= D_+ \Phi_1^+ = \bar{D}_- \Phi_2^-, & (10) \\ \mathcal{G} &= D_- \Phi_1^- = \bar{D}_+ \Phi_2^+, & (11) \end{aligned}$$

where the chiral Φ_p^+ and anti-chiral $\Phi_p^-, p = 1, 2$ superfields are defined as

$$\begin{aligned} \Phi_1^\pm &= q_1^\pm(z^\pm, \bar{z}^\pm) + \theta^\pm \zeta_1^\pm(z^\pm, \bar{z}^\pm) + \bar{\theta}^\pm \zeta_2^\pm(z^\pm, \bar{z}^\pm) + \theta^\pm \bar{\theta}^\pm q_2^\pm(z^\pm, \bar{z}^\pm), \\ \Phi_2^\pm &= p_1^\pm(z^\pm, \bar{z}^\pm) + \theta^\pm \xi_1^\pm(z^\pm, \bar{z}^\pm) + \bar{\theta}^\pm \xi_2^\pm(z^\pm, \bar{z}^\pm) + \theta^\pm \bar{\theta}^\pm p_2^\pm(z^\pm, \bar{z}^\pm). \end{aligned}$$

The second equality in (10) implies

$$\zeta_1^+ = \xi_2^-, \qquad q_2^+ = \partial_{\bar{z}} p_1^-, \qquad p_2^- = -\partial_z q_1^+, \qquad \partial_z \zeta_2^+ = -\partial_{\bar{z}} \xi_1^-, \tag{12}$$

whilst the second equality in (11) implies

$$\zeta_1^- = \xi_2^+, \qquad p_2^+ = -\partial_z q_1^-, \qquad q_2^- = \partial_{\bar{z}} p_1^+, \qquad \partial_z \zeta_2^- = -\partial_{\bar{z}} \xi_1^+. \tag{13}$$

Eqns. (3)-(9) describe the Backlund transformation for the $N = 2$ super sine-Gordon system. In appendix A we present these equations in components.

3 The Permutability condition

A Backlund transformation from $\phi_0^\pm$ to $\phi_1^\pm$ is described by

$$\begin{aligned}
D_+(\phi_0^+ - \phi_1^+) &= -\frac{8}{\kappa_1}\mathcal{F}^{(0,1)} \cos\left(\frac{\phi_0^- + \phi_1^-}{2}\right), && (14)\\
\bar{D}_+(\phi_0^+ + \phi_1^+) &= \kappa_1 \mathcal{G}^{(0,1)} \cos\left(\frac{\phi_0^- - \phi_1^-}{2}\right), && (15)\\
D_-(\phi_0^- - \phi_1^-) &= \lambda_1 \mathcal{G}^{(0,1)} \cos\left(\frac{\phi_0^+ + \phi_1^+}{2}\right), && (16)\\
\bar{D}_-(\phi_0^- + \phi_1^-) &= -\frac{8}{\lambda_1}\mathcal{F}^{(0,1)} \cos\left(\frac{\phi_0^+ - \phi_1^+}{2}\right), && (17)
\end{aligned}$$

where we have introduced the superscript indices $(0,1)$ for the auxiliary fermionic superfields denoting its dependence in $\phi_0^\pm$ and $\phi_1^\pm$. The later, in turn satisfy the following condition (as in (5) and (8))

$$\begin{aligned}
\bar{D}_+\mathcal{F}^{(0,1)} &= -g\frac{\kappa_1}{4} \sin\left(\frac{\phi_0^- - \phi_1^-}{2}\right), && (18)\\
D_+\mathcal{G}^{(0,1)} &= -g\frac{2}{\kappa_1} \sin\left(\frac{\phi_0^- + \phi_1^-}{2}\right), && (19)\\
\bar{D}_-\mathcal{G}^{(0,1)} &= g\frac{2}{\lambda_1} \sin\left(\frac{\phi_0^+ - \phi_1^+}{2}\right), && (20)\\
D_-\mathcal{F}^{(0,1)} &= g\frac{\lambda_1}{4} \sin\left(\frac{\phi_0^+ + \phi_1^+}{2}\right). && (21)
\end{aligned}$$

The chiral conditions (9), i.e.,

$$\bar{D}_-\mathcal{F}^{(0,1)} = 0, \qquad D_+\mathcal{F}^{(0,1)} = 0, \qquad \bar{D}_+\mathcal{G}^{(0,1)} = 0, \qquad D_-\mathcal{G}^{(0,1)} = 0,$$

are automatically satisfied using eqn. (10), i.e., expressing the fermionic superfields as derivatives of chiral superfields,

$$\mathcal{F}^{(0,1)} = D_+\Phi_1^{+(0,1)} = \bar{D}_-\Phi_2^{-(0,1)}, \qquad \mathcal{G}^{(0,1)} = D_-\Phi_1^{-(0,1)} = \bar{D}_+\Phi_2^{+(0,1)},$$

where the superscript indices indicate whether the superfield $\Phi_1^\pm$ and $\Phi_2^\pm$ depend upon $\phi_0^\pm$ and $\phi_1^\pm$. Acting with super derivatives $D_-, \bar{D}_-, D_+$ and $\bar{D}_+$ on eqns.(14), (15), (16) and (17) respectively, we find

$$\begin{aligned}
\partial_z(\phi_0^+ - \phi_1^+) &= -2\gamma_1\, s_{0,1}^+ c_{0,1}^- + \frac{8}{\kappa_1}\mathcal{F}^{(0,1)} D_- c_{0,1}^-, && (22)\\
\partial_{\bar{z}}(\phi_0^+ + \phi_1^+) &= 2\frac{g^2}{\gamma_1}\, \bar{s}_{0,1}^+ \bar{c}_{0,1}^- - \kappa_1 \mathcal{G}^{(0,1)} \bar{D}_- \bar{c}_{0,1}^-, && (23)\\
\partial_z(\phi_0^- - \phi_1^-) &= -2\gamma_1\, s_{0,1}^- c_{0,1}^+ - \lambda_1 \mathcal{G}^{(0,1)} D_+ c_{0,1}^+, && (24)\\
\partial_{\bar{z}}(\phi_0^- + \phi_1^-) &= 2\frac{g^2}{\gamma_1} \bar{s}_{0,1}^- \bar{c}_{0,1}^+ + \frac{8}{\lambda_1}\mathcal{F}^{(0,1)} \bar{D}_+ \bar{c}_{0,1}^+, && (25)
\end{aligned}$$

where $\gamma_1 = g\frac{\lambda_1}{\kappa_1}$ and

$$\begin{aligned} c^{\pm}_{j,k} &= \cos\left(\frac{\phi^{\pm}_j + \phi^{\pm}_k}{2}\right), \qquad s^{\pm}_{j,k} = \sin\left(\frac{\phi^{\pm}_j + \phi^{\pm}_k}{2}\right), && (26)\\ \bar{c}^{\pm}_{j,k} &= \cos\left(\frac{\phi^{\pm}_j - \phi^{\pm}_k}{2}\right), \qquad \bar{s}^{\pm}_{j,k} = \sin\left(\frac{\phi^{\pm}_j - \phi^{\pm}_k}{2}\right), && (27)\end{aligned}$$

We now assume that the order of two successive Backlund transformations is irrelevant leading to the same final result. Such condition is known as the permutability theorem, i.e., $\phi^{\pm}_0 \xrightarrow{\gamma_1} \phi^{\pm}_1 \xrightarrow{\gamma_2} \phi^{\pm}_{12}$ and in the inverse order, $\phi^{\pm}_0 \xrightarrow{\gamma_2} \phi^{\pm}_2 \xrightarrow{\gamma_1} \phi^{\pm}_{21}$, does not change the final result, $\phi^{\pm}_{12} = \phi^{\pm}_{21} \equiv \phi^{\pm}_3$.

The permutability theorem applied to the Backlund equation (14) leads to

$$\begin{aligned} D_+(\phi^+_0 - \phi^+_1) &= -\frac{8}{\kappa_1}\mathcal{F}^{(0,1)}c^-_{0,1},\\ D_+(\phi^+_1 - \phi^+_3) &= -\frac{8}{\kappa_2}\mathcal{F}^{(1,3)}c^-_{1,3},\\ D_+(\phi^+_0 - \phi^+_2) &= -\frac{8}{\kappa_2}\mathcal{F}^{(0,2)}c^-_{0,2},\\ D_+(\phi^+_2 - \phi^+_3) &= -\frac{8}{\kappa_1}\mathcal{F}^{(2,3)}c^-_{2,3}. && (28)\end{aligned}$$

Taking into account that the sum of the first two and the last two equations are the same, we obtain,

$$\frac{1}{\kappa_1}\mathcal{F}^{(0,1)}c^-_{0,1} + \frac{1}{\kappa_2}\mathcal{F}^{(1,3)}c^-_{1,3} = \frac{1}{\kappa_2}\mathcal{F}^{(0,2)}c^-_{0,2} + \frac{1}{\kappa_1}\mathcal{F}^{(2,3)}c^-_{2,3}. \tag{29}$$

Similarly, from (17), we obtain

$$\begin{aligned} \bar{D}_-(\phi^-_0 + \phi^-_1) &= -\frac{8}{\lambda_1}\mathcal{F}^{(0,1)}\bar{c}^+_{0,1},\\ \bar{D}_-(\phi^-_1 + \phi^-_3) &= -\frac{8}{\lambda_2}\mathcal{F}^{(1,3)}\bar{c}^+_{1,3},\\ \bar{D}_-(\phi^-_0 + \phi^-_2) &= -\frac{8}{\lambda_2}\mathcal{F}^{(0,2)}\bar{c}^+_{0,2},\\ \bar{D}_-(\phi^-_2 + \phi^-_3) &= -\frac{8}{\lambda_1}\mathcal{F}^{(2,3)}\bar{c}^+_{2,3}, && (30)\end{aligned}$$

leading to

$$\frac{1}{\lambda_1}\mathcal{F}^{(0,1)}\bar{c}^+_{0,1} - \frac{1}{\lambda_2}\mathcal{F}^{(1,3)}\bar{c}^+_{1,3} = \frac{1}{\lambda_2}\mathcal{F}^{(0,2)}\bar{c}^+_{0,2} - \frac{1}{\lambda_1}\mathcal{F}^{(2,3)}\bar{c}^+_{2,3}. \tag{31}$$

We propose as solution for the non-linear superposition formula $\phi^{\pm}_{12} = \phi^{\pm}_{21} = \phi^{\pm}_3$,

$$\phi^{\pm}_3 = \phi^{\pm}_0 + \Gamma_{\pm} + \Delta_{\pm}, \tag{32}$$

with

$$\begin{aligned}
&\Gamma_{\pm}(x,y) = 2\arctan\left[\delta\tan\left(\frac{x+y}{4}\right)\right] \pm 2\arctan\left[\delta\tan\left(\frac{x-y}{4}\right)\right], \\
&x = \phi_1^+ - \phi_2^+, \qquad y = \phi_1^- - \phi_2^-, \\
&\delta = \frac{\gamma_1+\gamma_2}{\gamma_1-\gamma_2}, \qquad \gamma_k = g\frac{\lambda_k}{\kappa_k}.
\end{aligned} \tag{33}$$

Notice that the solution $\phi_3^{\pm}$ when the fermionic superfields are neglected is derived in the appendix B to be $\phi_3^{\pm} = \phi_0^{\pm} + \Gamma_{\pm}$. The term $\Delta_{\pm}$ comes from the contribution of the fermionic superfields and has the following form

$$\begin{aligned}
&\Delta_{\pm} = \sum_{j,k=1}^{2} \Lambda_{j,k}^{\pm} f_{j,k} + \Lambda_0^{\pm} f_0, \\
&f_{j,k} = \mathcal{F}^{(0,j)}\mathcal{G}^{(0,k)}, \qquad f_0 = \mathcal{F}^{(0,1)}\mathcal{F}^{(0,2)}\mathcal{G}^{(0,1)}\mathcal{G}^{(0,2)},
\end{aligned}$$

where we have assumed the coefficients $\Lambda^{\pm}$ to be functionals of $x = (\phi_1^+ - \phi_2^+)$ and $y = (\phi_1^- - \phi_2^-)$,i.e.,

$$\Lambda_{j,k}^{\pm} = \Lambda_{j,k}^{\pm}(x,y), \qquad \Lambda_0^{\pm} = \Lambda_0^{\pm}(x,y). \tag{34}$$

Observe that there are no terms like $\Lambda_1^{\pm}\mathcal{F}^{(0,1)}\mathcal{F}^{(0,2)}$ nor $\Lambda_2^{\pm}\mathcal{G}^{(0,1)}\mathcal{G}^{(0,2)}$ due to chiral equations (2). $\Lambda_{\pm}$ are determined in appendix C where,

$$\begin{aligned}
\Lambda_{1,1}^{+} &= \Lambda_{2,2}^{+} = -\frac{8\mu_-}{g\eta_+\eta_-}\cos\left(\frac{x}{2}\right)\sin\left(\frac{y}{2}\right), \\
\Lambda_{1,2}^{+} &= \frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\lambda_2}{\lambda_1}\right)\sin\left(\frac{y}{2}\right), \\
\Lambda_{2,1}^{+} &= \frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\lambda_1}{\lambda_2}\right)\sin\left(\frac{y}{2}\right), \\
\Lambda_{0}^{+} &= -\frac{32\mu_-}{(g\eta_+\eta_-)^2}\sin\left(\frac{x}{2}\right)\left[\cos\left(\frac{y}{2}\right)(a+\cos x-\cos y) - 2\mu_+\cos\left(\frac{x}{2}\right)\right], \\
\\
\Lambda_{1,1}^{-} &= \Lambda_{2,2}^{-} = \frac{8\mu_-}{g\eta_+\eta_-}\cos\left(\frac{y}{2}\right)\sin\left(\frac{x}{2}\right), \\
\Lambda_{1,2}^{-} &= -\frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\kappa_2}{\kappa_1}\right)\sin\left(\frac{x}{2}\right), \\
\Lambda_{2,1}^{-} &= -\frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\kappa_1}{\kappa_2}\right)\sin\left(\frac{x}{2}\right), \\
\Lambda_{0}^{-} &= -\frac{32\mu_-}{(g\eta_+\eta_-)^2}\sin\left(\frac{y}{2}\right)\left[\cos\left(\frac{x}{2}\right)(a-\cos x+\cos y) - 2\mu_+\cos\left(\frac{y}{2}\right)\right],
\end{aligned}$$

$$\begin{aligned}
\mu_{\pm} &= \frac{\gamma_1}{\gamma_2} \pm \frac{\gamma_2}{\gamma_1}, \\
a &= \frac{1}{2}\left(\frac{\gamma_1^2}{\gamma_2^2} + \frac{\gamma_2^2}{\gamma_1^2}\right) + 3, \\
\eta_{\pm} &= \mu_+ - 2\cos\left(\frac{x \pm y}{2}\right).
\end{aligned}$$

3.1 Solution in Components

In components the non-linear superposition formula (32) yields the following expressions,

$$\begin{aligned}
\varphi_3^+ &= \varphi_0^+ + \tilde{\Gamma}_+ - \frac{8\mu_-}{g\tilde{\eta}_+\tilde{\eta}_-}\left(\mathcal{A}_1^+ - \mathcal{B}_1^+ + \frac{4}{g\tilde{\eta}_+\tilde{\eta}_-}\mathcal{C}_1^+\right), \\
\psi_3^- &= \psi_0^- + F_{1,2}\psi_{1,2}^- + \frac{8\mu_-}{g\tilde{\eta}_+\tilde{\eta}_-}\left(\mathcal{A}_2^+ - \mathcal{B}_2^+ + \frac{4}{g\tilde{\eta}_+\tilde{\eta}_-}\mathcal{C}_2^+\right), \\
\bar{\psi}_3^- &= \bar{\psi}_0^- + F_{1,2}\bar{\psi}_{1,2}^- + \frac{8\mu_-}{g\tilde{\eta}_+\tilde{\eta}_-}\left(\mathcal{A}_3^+ - \mathcal{B}_3^+ + \frac{4}{g\tilde{\eta}_+\tilde{\eta}_-}\mathcal{C}_3^+\right),
\end{aligned}$$

$$\begin{aligned}
\varphi_3^- &= \varphi_0^- + \tilde{\Gamma}_- + \frac{8\mu_-}{g\tilde{\eta}_+\tilde{\eta}_-}\left(\mathcal{A}_1^- - \mathcal{B}_1^- - \frac{4}{g\tilde{\eta}_+\tilde{\eta}_-}\mathcal{C}_1^-\right), \\
\psi_3^+ &= \psi_0^+ + F_{1,2}\psi_{1,2}^+ - \frac{8\mu_-}{g\tilde{\eta}_+\tilde{\eta}_-}\left(\mathcal{A}_2^- - \mathcal{B}_2^- - \frac{4}{g\tilde{\eta}_+\tilde{\eta}_-}\mathcal{C}_2^-\right), \\
\bar{\psi}_3^+ &= \bar{\psi}_0^+ + F_{1,2}\bar{\psi}_{1,2}^+ - \frac{8\mu_-}{g\tilde{\eta}_+\tilde{\eta}_-}\left(\mathcal{A}_3^- - \mathcal{B}_3^- - \frac{4}{g\tilde{\eta}_+\tilde{\eta}_-}\mathcal{C}_3^-\right),
\end{aligned}$$

where

$$\begin{aligned}
\tilde{\Gamma}_\pm &= 2\arctan\left[\delta\tan\left(\frac{\varphi_{1,2}^+ + \varphi_{1,2}^-}{4}\right)\right] \pm 2\arctan\left[\delta\tan\left(\frac{\varphi_{1,2}^+ - \varphi_{1,2}^-}{4}\right)\right], \\
\tilde{\eta}_\pm &= \mu_+ - 2\cos\left(\frac{\varphi_{1,2}^+ \pm \varphi_{1,2}^-}{2}\right), \\
F_{1,2} &= \frac{\delta}{2}\left[\frac{\sec^2\left(\frac{\varphi_{1,2}^+ + \varphi_{1,2}^-}{4}\right)}{1+\delta^2\tan^2\left(\frac{\varphi_{1,2}^+ + \varphi_{1,2}^-}{4}\right)} + \frac{\sec^2\left(\frac{\varphi_{1,2}^+ - \varphi_{1,2}^-}{4}\right)}{1+\delta^2\tan^2\left(\frac{\varphi_{1,2}^+ - \varphi_{1,2}^-}{4}\right)}\right],
\end{aligned}$$

$$
\begin{aligned}
\mathcal{A}_1^+ &= \cos\left(\frac{\varphi_{1,2}^+}{2}\right)\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\left(\zeta_1^{+(0,1)}\xi_2^{+(0,1)}+\zeta_1^{+(0,2)}\xi_2^{+(0,2)}\right),\\
\mathcal{A}_2^+ &= -\cos\left(\frac{\varphi_{1,2}^+}{2}\right)\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\left(\zeta_1^{+(0,1)}p_2^{+(0,1)}+\zeta_1^{+(0,2)}p_2^{+(0,2)}\right)\\
&\quad -\Sigma^+\left(\zeta_1^{+(0,1)}\xi_2^{+(0,1)}+\zeta_1^{+(0,2)}\xi_2^{+(0,2)}\right)\psi_{1,2}^-,\\
\mathcal{A}_3^+ &= -\cos\left(\frac{\varphi_{1,2}^+}{2}\right)\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\left(q_2^{+(0,1)}\xi_2^{+(0,1)}+q_2^{+(0,2)}\xi_2^{+(0,2)}\right)\\
&\quad -\Sigma^+\left(\zeta_1^{+(0,1)}\xi_2^{+(0,1)}+\zeta_1^{+(0,2)}\xi_2^{+(0,2)}\right)\bar{\psi}_{1,2}^-,
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{B}_1^+ &= \sin\left(\frac{\varphi_{1,2}^-}{2}\right)\left(\frac{\lambda_2}{\lambda_1}\zeta_1^{+(0,1)}\xi_2^{+(0,2)}+\frac{\lambda_1}{\lambda_2}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\right),\\
\mathcal{B}_2^+ &= -\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\left(\frac{\lambda_2}{\lambda_1}\zeta_1^{+(0,1)}p_2^{+(0,2)}+\frac{\lambda_1}{\lambda_2}\zeta_1^{+(0,2)}p_2^{+(0,1)}\right)\\
&\quad -\Omega^+\left(\frac{\lambda_2}{\lambda_1}\zeta_1^{+(0,1)}\xi_2^{+(0,2)}+\frac{\lambda_1}{\lambda_2}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\right)\psi_{1,2}^-,\\
\mathcal{B}_3^+ &= -\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\left(\frac{\lambda_2}{\lambda_1}q_2^{+(0,1)}\xi_2^{+(0,2)}+\frac{\lambda_1}{\lambda_2}q_2^{+(0,2)}\xi_2^{+(0,1)}\right)\\
&\quad -\Omega^+\left(\frac{\lambda_2}{\lambda_1}\zeta_1^{+(0,1)}\xi_2^{+(0,2)}+\frac{\lambda_1}{\lambda_2}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\right)\bar{\psi}_{1,2}^-,
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{C}_1^+ &= \sin\left(\frac{\varphi_{1,2}^+}{2}\right)A^+\zeta_1^{+(0,1)}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\xi_2^{+(0,2)},\\
\mathcal{C}_2^+ &= \sin\left(\frac{\varphi_{1,2}^+}{2}\right)A^+\zeta_1^{+(0,1)}\zeta_1^{+(0,2)}\left(\xi_2^{+(0,2)}p_2^{+(0,1)}-\xi_2^{+(0,1)}p_2^{+(0,2)}\right),\\
\mathcal{C}_3^+ &= \sin\left(\frac{\varphi_{1,2}^+}{2}\right)A^+\xi_2^{+(0,1)}\xi_2^{+(0,2)}\left(\zeta_1^{+(0,1)}q_2^{+(0,2)}-\zeta_1^{+(0,2)}q_2^{+(0,1)}\right),
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{A}_1^- &= \cos\left(\frac{\varphi_{1,2}^-}{2}\right)\sin\left(\frac{\varphi_{1,2}^+}{2}\right)\left(\zeta_1^{+(0,1)}\xi_2^{+(0,1)}+\zeta_1^{+(0,2)}\xi_2^{+(0,2)}\right),\\
\mathcal{A}_2^- &= -\cos\left(\frac{\varphi_{1,2}^-}{2}\right)\sin\left(\frac{\varphi_{1,2}^+}{2}\right)\left(\partial_z q_1^{+(0,1)}\xi_2^{+(0,1)}+\partial_z q_1^{+(0,2)}\xi_2^{+(0,2)}\right)\\
&\quad -\Sigma^-\left(\zeta_1^{+(0,1)}\xi_2^{+(0,1)}+\zeta_1^{+(0,2)}\xi_2^{+(0,2)}\right)\psi_{1,2}^+,\\
\mathcal{A}_3^- &= \cos\left(\frac{\varphi_{1,2}^-}{2}\right)\sin\left(\frac{\varphi_{1,2}^+}{2}\right)\left(\zeta_1^{+(0,1)}\partial_{\bar{z}} p_1^{+(0,1)}+\zeta_1^{+(0,2)}\partial_{\bar{z}} p_1^{+(0,2)}\right)\\
&\quad -\Sigma^-\left(\zeta_1^{+(0,1)}\xi_2^{+(0,1)}+\zeta_1^{+(0,2)}\xi_2^{+(0,2)}\right)\bar{\psi}_{1,2}^+,
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{B}_1^- &= \sin\left(\frac{\varphi_{1,2}^+}{2}\right)\left(\frac{\kappa_2}{\kappa_1}\zeta_1^{+(0,1)}\xi_2^{+(0,2)}+\frac{\kappa_1}{\kappa_2}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\right),\\
\mathcal{B}_2^- &= -\sin\left(\frac{\varphi_{1,2}^+}{2}\right)\left(\frac{\kappa_2}{\kappa_1}\partial_z q_1^{+(0,1)}\xi_2^{+(0,2)}+\frac{\kappa_1}{\kappa_2}\partial_z q_1^{+(0,2)}\xi_2^{+(0,1)}\right)\\
&\quad -\Omega^-\left(\frac{\kappa_2}{\kappa_1}\zeta_1^{+(0,1)}\xi_2^{+(0,2)}+\frac{\kappa_1}{\kappa_2}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\right)\psi_{1,2}^+,\\
\mathcal{B}_3^- &= \sin\left(\frac{\varphi_{1,2}^+}{2}\right)\left(\frac{\kappa_2}{\kappa_1}\zeta_1^{+(0,1)}\partial_{\bar{z}} p_1^{+(0,2)}+\frac{\kappa_1}{\kappa_2}\zeta_1^{+(0,2)}\partial_{\bar{z}} p_1^{+(0,1)}\right)\\
&\quad -\Omega^-\left(\frac{\kappa_2}{\kappa_1}\zeta_1^{+(0,1)}\xi_2^{+(0,2)}+\frac{\kappa_1}{\kappa_2}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\right)\bar{\psi}_{1,2}^+,
\end{aligned}
$$

$$
\begin{aligned}
\mathcal{C}_1^- &= \sin\left(\frac{\varphi_{1,2}^-}{2}\right)A^-\zeta_1^{+(0,1)}\zeta_1^{+(0,2)}\xi_2^{+(0,1)}\xi_2^{+(0,2)},\\
\mathcal{C}_2^- &= -\sin\left(\frac{\varphi_{1,2}^-}{2}\right)A^-\xi_2^{+(0,1)}\xi_2^{+(0,2)}\left(\zeta_1^{+(0,2)}\partial_z q_1^{+(0,1)}-\zeta_1^{+(0,1)}\partial_z q_1^{+(0,2)}\right),\\
\mathcal{C}_3^- &= -\sin\left(\frac{\varphi_{1,2}^-}{2}\right)A^-\zeta_1^{+(0,1)}\zeta_1^{+(0,2)}\left(\xi_2^{+(0,2)}\partial_{\bar{z}} p_1^{+(0,1)}-\xi_2^{+(0,1)}\partial_{\bar{z}} p_1^{+(0,2)}\right),
\end{aligned}
$$

$$\begin{aligned}
\Sigma^+ &= \cos\left(\frac{\varphi^+_{1,2}}{2}\right)\Omega^+ - \frac{1}{2}\sin\left(\frac{\varphi^-_{1,2}}{2}\right)\sin\left(\frac{\varphi^+_{1,2}}{2}\right),\\
\Omega^+ &= -\sin\left(\frac{\varphi^-_{1,2}}{2}\right)\left[\frac{1}{\tilde\eta_+}\sin\left(\frac{\varphi^+_{1,2}+\varphi^-_{1,2}}{2}\right) + \frac{1}{\tilde\eta_-}\sin\left(\frac{\varphi^+_{1,2}-\varphi^-_{1,2}}{2}\right)\right],\\
A^+ &= \cos\left(\frac{\varphi^-_{1,2}}{2}\right)\left(a + \cos\varphi^+_{1,2} - \cos\varphi^-_{1,2}\right) - 2\mu_+\cos\left(\frac{\varphi^+_{1,2}}{2}\right),
\end{aligned}$$

$$\begin{aligned}
\Sigma^- &= \cos\left(\frac{\varphi^-_{1,2}}{2}\right)\Omega^- - \frac{1}{2}\sin\left(\frac{\varphi^-_{1,2}}{2}\right)\sin\left(\frac{\varphi^+_{1,2}}{2}\right),\\
\Omega^- &= -\sin\left(\frac{\varphi^+_{1,2}}{2}\right)\left[\frac{1}{\tilde\eta_+}\sin\left(\frac{\varphi^+_{1,2}+\varphi^-_{1,2}}{2}\right) - \frac{1}{\tilde\eta_-}\sin\left(\frac{\varphi^+_{1,2}-\varphi^-_{1,2}}{2}\right)\right],\\
A^- &= \cos\left(\frac{\varphi^+_{1,2}}{2}\right)\left(a - \cos\varphi^+_{1,2} + \cos\varphi^-_{1,2}\right) - 2\mu_+\cos\left(\frac{\varphi^-_{1,2}}{2}\right),
\end{aligned}$$

and denoted

$$\varphi^\pm_{1,2} = \varphi^\pm_1 - \varphi^\pm_2, \qquad \psi^\pm_{1,2} = \psi^\pm_1 - \psi^\pm_2, \qquad \bar\psi^\pm_{1,2} = \bar\psi^\pm_1 - \bar\psi^\pm_2.$$

From the Backlund eqns. we get (see app. A)

$$\begin{aligned}
\zeta_1^{+(0,k)} &= -\frac{\kappa_k}{8}\frac{(\psi_0^- - \psi_k^-)}{\cos\left(\frac{\varphi_0^-+\varphi_k^-}{2}\right)}, & \xi_2^{+(0,k)} &= \frac{1}{\kappa_k}\frac{(\bar\psi_0^- + \bar\psi_k^-)}{\cos\left(\frac{\varphi_0^--\varphi_k^-}{2}\right)},\\
\partial_z q_1^{+(0,k)} &= \frac{\lambda_k g}{4}\sin\left(\frac{\varphi_0^+ + \varphi_k^+}{2}\right), & p_2^{+(0,k)} &= \frac{2g}{\kappa_k}\sin\left(\frac{\varphi_0^- + \varphi_k^-}{2}\right),\\
q_2^{+(0,k)} &= -\frac{\kappa_k g}{4}\sin\left(\frac{\varphi_0^- - \varphi_k^-}{2}\right), & \partial_{\bar z} p_1^{+(0,k)} &= \frac{2g}{\lambda_k}\sin\left(\frac{\varphi_0^+ - \varphi_k^+}{2}\right).
\end{aligned}$$

4 1-Soliton Solution

Setting $\phi_0^\pm = 0$ in the Backlund eqns. (14)-(21) we find in components,

$$\begin{aligned}
\partial_{\bar z}\zeta_1^{+(0,1)} &= -\frac{g^2}{\gamma_1}\cos\left(\frac{\varphi_1^+}{2}\right)\cos\left(\frac{\varphi_1^-}{2}\right)\zeta_1^{+(0,1)},\\
\partial_z\zeta_1^{+(0,1)} &= \gamma_1\cos\left(\frac{\varphi_1^+}{2}\right)\cos\left(\frac{\varphi_1^-}{2}\right)\zeta_1^{+(0,1)},\\
\partial_{\bar z}\xi_2^{+(0,1)} &= -\frac{g^2}{\gamma_1}\cos\left(\frac{\varphi_1^+}{2}\right)\cos\left(\frac{\varphi_1^-}{2}\right)\xi_2^{+(0,1)},\\
\partial_z\xi_2^{+(0,1)} &= \gamma_1\cos\left(\frac{\varphi_1^+}{2}\right)\cos\left(\frac{\varphi_1^-}{2}\right)\xi_2^{+(0,1)},
\end{aligned}$$

$$\begin{aligned}
\partial_{\bar z}\varphi_1^{\pm} &= -\frac{2g^2}{\gamma_1}\sin\left(\frac{\varphi_1^{\pm}}{2}\right)\cos\left(\frac{\varphi_1^{\mp}}{2}\right),\\
\partial_{z}\varphi_1^{\pm} &= 2\gamma_1\sin\left(\frac{\varphi_1^{\pm}}{2}\right)\cos\left(\frac{\varphi_1^{\mp}}{2}\right).
\end{aligned}$$

Integrating the above eqns. we get the 1-soliton solution,

$$\begin{aligned}
\psi_1^- &= \frac{8}{\kappa_1}\cos\left(\frac{\varphi_1^-}{2}\right)\zeta_1^{+(0,1)}, \qquad \psi_1^+ = -\lambda_1\cos\left(\frac{\varphi_1^+}{2}\right)\xi_2^{+(0,1)},\\
\bar\psi_1^- &= \kappa_1\cos\left(\frac{\varphi_1^-}{2}\right)\xi_2^{+(0,1)}, \qquad \bar\psi_1^+ = -\frac{8}{\lambda_1}\cos\left(\frac{\varphi_1^+}{2}\right)\zeta_1^{+(0,1)},
\end{aligned}$$

$$\begin{aligned}
\varphi_1^{\pm} &= 2\arctan(a_1\rho_1) \pm 2\arctan(b_1\rho_1),\\
\zeta_1^{+(0,1)} &= \xi_2^{+(0,1)} = \epsilon_1\chi_1,\\
\chi_1 &= \frac{\rho_1}{\sqrt{(1+a_1^2\rho_1^2)(1+b_1^2\rho_1^2)}},
\end{aligned}$$

where a_1 and b_1 are arbitrary constants, ϵ_1 is a grassmann parameter and

$$\rho_1 = \exp\left(\gamma_1 z - \frac{g^2}{\gamma_1}\bar z\right).$$

The 1-soliton solution constructed in this section can be obtained from those of [7] by relating parameters since they both involve a single grassmann parameter.

5 2-Soliton Solution

For the 2-soliton case we obtain from the superposition formulae (32)

$$\begin{aligned}
\varphi_3^+ &= \varphi_3^{+(0)} + \varphi_3^{+(1)}\epsilon_1\epsilon_2,\\
\varphi_3^{+(0)} &= 2\arctan\left[\delta\tan\left(\frac{\varphi_{1,2}^+ + \varphi_{1,2}^-}{4}\right)\right] + 2\arctan\left[\delta\tan\left(\frac{\varphi_{1,2}^+ - \varphi_{1,2}^-}{4}\right)\right],\\
\varphi_3^{+(1)} &= \frac{8\mu_-}{g\tilde\eta_+\tilde\eta_-}\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\left(\frac{\lambda_2}{\lambda_1}-\frac{\lambda_1}{\lambda_2}\right)\chi_1\chi_2,
\end{aligned}$$

$$\begin{aligned}
\psi_3^- &= \epsilon_1\psi_3^{-(1)} + \epsilon_2\psi_3^{-(2)},\\
\psi_3^{-(1)} &= \frac{8}{\kappa_1}F_{1,2}\cos\left(\frac{\varphi_1^-}{2}\right)\chi_1\\
&\quad +\frac{16}{\kappa_1\gamma_1}\frac{\mu_-}{\tilde\eta_+\tilde\eta_-}\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\chi_1\left[\gamma_2\sin\left(\frac{\varphi_2^-}{2}\right) - \gamma_1\cos\left(\frac{\varphi_{1,2}^+}{2}\right)\sin\left(\frac{\varphi_1^-}{2}\right)\right],\\
\psi_3^{-(2)} &= -\frac{8}{\kappa_2}F_{1,2}\cos\left(\frac{\varphi_2^-}{2}\right)\chi_2\\
&\quad +\frac{16}{\kappa_2\gamma_2}\frac{\mu_-}{\tilde\eta_+\tilde\eta_-}\sin\left(\frac{\varphi_{1,2}^-}{2}\right)\chi_2\left[\gamma_1\sin\left(\frac{\varphi_1^-}{2}\right) - \gamma_2\cos\left(\frac{\varphi_{1,2}^+}{2}\right)\sin\left(\frac{\varphi_2^-}{2}\right)\right],
\end{aligned}$$

$$\begin{aligned}
\bar{\psi}_3^- &= \epsilon_1 \bar{\psi}_3^{-(1)} + \epsilon_2 \bar{\psi}_3^{-(2)}, \\
\bar{\psi}_3^{-(1)} &= \kappa_1 F_{1,2} \cos\left(\frac{\varphi_1^-}{2}\right) \chi_1 \\
&\quad + \frac{2\kappa_1}{\gamma_2} \frac{\mu_-}{\tilde{\eta}_+ \tilde{\eta}_-} \sin\left(\frac{\varphi_{1,2}^-}{2}\right) \chi_1 \left[\gamma_1 \sin\left(\frac{\varphi_2^-}{2}\right) - \gamma_2 \cos\left(\frac{\varphi_{1,2}^+}{2}\right) \sin\left(\frac{\varphi_1^-}{2}\right)\right], \\
\bar{\psi}_3^{-(2)} &= -\kappa_2 F_{1,2} \cos\left(\frac{\varphi_2^-}{2}\right) \chi_2 \\
&\quad + \frac{2\kappa_2}{\gamma_1} \frac{\mu_-}{\tilde{\eta}_+ \tilde{\eta}_-} \sin\left(\frac{\varphi_{1,2}^-}{2}\right) \chi_2 \left[\gamma_2 \sin\left(\frac{\varphi_1^-}{2}\right) - \gamma_1 \cos\left(\frac{\varphi_{1,2}^+}{2}\right) \sin\left(\frac{\varphi_2^-}{2}\right)\right],
\end{aligned}$$

$$\begin{aligned}
\varphi_3^- &= \varphi_3^{-(0)} + \varphi_3^{-(1)} \epsilon_1 \epsilon_2, \\
\varphi_3^{-(0)} &= 2 \arctan\left[\delta \tan\left(\frac{\varphi_{1,2}^+ + \varphi_{1,2}^-}{4}\right)\right] - 2 \arctan\left[\delta \tan\left(\frac{\varphi_{1,2}^+ - \varphi_{1,2}^-}{4}\right)\right], \\
\varphi_3^{-(1)} &= -\frac{8\mu_-}{g \tilde{\eta}_+ \tilde{\eta}_-} \sin\left(\frac{\varphi_{1,2}^+}{2}\right) \left(\frac{\kappa_2}{\kappa_1} - \frac{\kappa_1}{\kappa_2}\right) \chi_1 \chi_2,
\end{aligned}$$

$$\begin{aligned}
\psi_3^+ &= \epsilon_1 \psi_3^{+(1)} + \epsilon_2 \psi_3^{+(2)}, \\
\psi_3^{+(1)} &= -\lambda_1 F_{1,2} \cos\left(\frac{\varphi_1^+}{2}\right) \chi_1 \\
&\quad - \frac{2\lambda_1}{\gamma_1} \frac{\mu_-}{\tilde{\eta}_+ \tilde{\eta}_-} \sin\left(\frac{\varphi_{1,2}^+}{2}\right) \chi_1 \left[\gamma_2 \sin\left(\frac{\varphi_2^+}{2}\right) - \gamma_1 \cos\left(\frac{\varphi_{1,2}^-}{2}\right) \sin\left(\frac{\varphi_1^+}{2}\right)\right], \\
\psi_3^{+(2)} &= \lambda_2 F_{1,2} \cos\left(\frac{\varphi_2^+}{2}\right) \chi_2 \\
&\quad - \frac{2\lambda_2}{\gamma_2} \frac{\mu_-}{\tilde{\eta}_+ \tilde{\eta}_-} \sin\left(\frac{\varphi_{1,2}^+}{2}\right) \chi_2 \left[\gamma_1 \sin\left(\frac{\varphi_1^+}{2}\right) - \gamma_2 \cos\left(\frac{\varphi_{1,2}^-}{2}\right) \sin\left(\frac{\varphi_2^+}{2}\right)\right],
\end{aligned}$$

$$\begin{aligned}
\bar{\psi}_3^+ &= \epsilon_1 \bar{\psi}_3^{+(1)} + \epsilon_2 \bar{\psi}_3^{+(2)}, \\
\bar{\psi}_3^{+(1)} &= -\frac{8}{\lambda_1} F_{1,2} \cos\left(\frac{\varphi_1^+}{2}\right) \chi_1 \\
&\quad - \frac{16}{\lambda_1 \gamma_2} \frac{\mu_-}{\tilde{\eta}_+ \tilde{\eta}_-} \sin\left(\frac{\varphi_{1,2}^+}{2}\right) \chi_1 \left[\gamma_1 \sin\left(\frac{\varphi_2^+}{2}\right) - \gamma_2 \cos\left(\frac{\varphi_{1,2}^-}{2}\right) \sin\left(\frac{\varphi_1^+}{2}\right)\right], \\
\bar{\psi}_3^{+(2)} &= \frac{8}{\lambda_2} F_{1,2} \cos\left(\frac{\varphi_2^+}{2}\right) \chi_2 \\
&\quad - \frac{16}{\lambda_2 \gamma_1} \frac{\mu_-}{\tilde{\eta}_+ \tilde{\eta}_-} \sin\left(\frac{\varphi_{1,2}^+}{2}\right) \chi_2 \left[\gamma_2 \sin\left(\frac{\varphi_1^+}{2}\right) - \gamma_1 \cos\left(\frac{\varphi_{1,2}^-}{2}\right) \sin\left(\frac{\varphi_2^+}{2}\right)\right],
\end{aligned}$$

where

$$
\begin{aligned}
\varphi_k^{\pm} &= 2\arctan(a_k\rho_k) \pm 2\arctan(b_k\rho_k), \\
\chi_k &= \frac{\rho_k}{\sqrt{(1+a_k^2\rho_k^2)(1+b_k^2\rho_k^2)}},
\end{aligned}
$$

$k = 1, 2$, a_k and b_k are arbitrary constants, ϵ_k is a grassmann constant and

$$\rho_k = \exp\left(\gamma_k z - \frac{g^2}{\gamma_k}\bar{z}\right).$$

Notice that the 2-soliton solution constructed in this section generalizes those constructed in ref. [7]) involving a single grassmann parameter.

Both 1- and 2-soliton solutions presented above were verified to satisfy the equations of motion.

Acknowledgments

LHY acknowledges support from Fapesp, JFG and AHZ thank CNPq for partial support.

Appendix A

In order to simplify notation let us introduce $\varphi_{\pm}^{(-)} = \varphi_1^- \pm \varphi_2^-$, $\varphi_{\pm}^{(+)} = \varphi_1^+ \pm \varphi_2^+$ and similar notation for the other fields.

In components eqn. (5) becomes

- $\bar{D}_+\mathcal{F} = -\frac{\kappa g}{4}\sin\left(\frac{\phi_1^- - \phi_2^-}{2}\right)$,

$$\Downarrow$$

$$q_2^+ = -\frac{\kappa g}{4}\sin\left(\frac{\varphi_-^{(-)}}{2}\right), \qquad \partial_{\bar{z}}\zeta_1^+ = -\frac{\kappa g}{8}\cos\left(\frac{\varphi_-^{(-)}}{2}\right)\bar{\psi}_-^{(+)}, \qquad \partial_z\zeta_2^+ = \frac{\kappa g}{8}\cos\left(\frac{\varphi_-^{(-)}}{2}\right)\psi_-^{(+)},$$

$$\partial_{\bar{z}}\partial_z q_1^+ = \frac{\kappa g}{8}\cos\left(\frac{\varphi_-^{(-)}}{2}\right)F_-^{(-)} + \frac{\kappa g}{16}\sin\left(\frac{\varphi_-^{(-)}}{2}\right)\psi_-^{(+)}\bar{\psi}_-^{(+)}.$$

- $D_+\mathcal{G} = -\frac{2g}{\kappa}\sin\left(\frac{\phi_1^- + \phi_2^-}{2}\right)$,

$$\Downarrow$$

$$p_2^+ = \frac{2g}{\kappa}\sin\left(\frac{\varphi_+^{(-)}}{2}\right), \qquad \partial_{\bar{z}}\xi_1^+ = \frac{g}{\kappa}\cos\left(\frac{\varphi_+^{(-)}}{2}\right)\bar{\psi}_+^{(+)}, \qquad \partial_z\xi_2^+ = -\frac{g}{\kappa}\cos\left(\frac{\varphi_+^{(-)}}{2}\right)\psi_+^{(+)},$$

$$\partial_{\bar{z}}\partial_z p_1^+ = -\frac{g}{\kappa}\cos\left(\frac{\varphi_+^{(-)}}{2}\right)F_+^{(-)} - \frac{g}{2\kappa}\sin\left(\frac{\varphi_+^{(-)}}{2}\right)\psi_+^{(+)}\bar{\psi}_+^{(+)}.$$

Similarly we find for (8),

- $\bar{D}_-\mathcal{G} = \frac{2g}{\lambda}\sin\left(\frac{\phi_1^+ - \phi_2^+}{2}\right),$

$$\Downarrow$$

$$q_2^- = \frac{2g}{\lambda}\sin\left(\frac{\varphi_-^{(+)}}{2}\right), \qquad \partial_{\bar{z}}\zeta_1^- = \frac{g}{\lambda}\cos\left(\frac{\varphi_-^{(+)}}{2}\right)\bar{\psi}_-^{(-)}, \qquad \partial_z\zeta_2^- = -\frac{g}{\lambda}\cos\left(\frac{\varphi_-^{(+)}}{2}\right)\psi_-^{(-)},$$

$$\partial_{\bar{z}}\partial_z q_1^- = -\frac{g}{\lambda}\cos\left(\frac{\varphi_-^{(+)}}{2}\right)F_-^{(+)} - \frac{g}{2\lambda}\sin\left(\frac{\varphi_-^{(+)}}{2}\right)\psi_-^{(-)}\bar{\psi}_-^{(-)}.$$

- $D_-\mathcal{F} = \frac{\lambda g}{4}\sin\left(\frac{\phi_1^+ + \phi_2^+}{2}\right),$

$$\Downarrow$$

$$p_2^- = -\frac{\lambda g}{4}\sin\left(\frac{\varphi_+^{(+)}}{2}\right), \qquad \partial_{\bar{z}}\xi_1^- = -\frac{\lambda g}{8}\cos\left(\frac{\varphi_+^{(+)}}{2}\right)\bar{\psi}_+^{(-)}, \qquad \partial_z\xi_2^- = \frac{\lambda g}{8}\cos\left(\frac{\varphi_+^{(+)}}{2}\right)\psi_+^{(-)},$$

$$\partial_{\bar{z}}\partial_z p_1^- = \frac{\lambda g}{8}\cos\left(\frac{\varphi_+^{(+)}}{2}\right)F_+^{(+)} + \frac{\lambda g}{16}\sin\left(\frac{\varphi_+^{(+)}}{2}\right)\psi_+^{(-)}\bar{\psi}_+^{(-)}.$$

From (3) and (4),

- $D_+\phi_1^+ = D_+\phi_2^+ - \frac{8}{\kappa}\mathcal{F}\cos\left(\frac{\phi_1^- + \phi_2^-}{2}\right),$

$$\Downarrow$$

$$\psi_-^{(-)} = -\frac{8}{\kappa}\zeta_1^+\cos\left(\frac{\varphi_+^{(-)}}{2}\right), \qquad F_-^{(+)} = -\frac{8}{\kappa}q_2^+\cos\left(\frac{\varphi_+^{(-)}}{2}\right),$$

$$\partial_z\varphi_-^{(+)} = -\frac{4}{\kappa}\sin\left(\frac{\varphi_+^{(-)}}{2}\right)\zeta_1^+\psi_+^{(+)} - \frac{8}{\kappa}\partial_z q_1^+\cos\left(\frac{\varphi_+^{(-)}}{2}\right).$$

- $\bar{D}_+\phi_1^+ = -\bar{D}_+\phi_2^+ + \kappa\mathcal{G}\cos\left(\frac{\phi_1^- - \phi_2^-}{2}\right),$

$$\Downarrow$$

$$\bar{\psi}_+^{(-)} = \kappa\xi_2^+\cos\left(\frac{\varphi_-^{(-)}}{2}\right), \qquad F_+^{(+)} = \kappa p_2^+\cos\left(\frac{\varphi_-^{(-)}}{2}\right),$$

$$\partial_{\bar{z}}\varphi_+^{(+)} = \frac{\kappa}{2}\sin\left(\frac{\varphi_-^{(-)}}{2}\right)\xi_2^+\bar{\psi}_-^{(+)} + \kappa\partial_{\bar{z}}p_1^+\cos\left(\frac{\varphi_-^{(-)}}{2}\right).$$

From (6) and (7),

- $D_-\phi_1^- = D_-\phi_2^- + \lambda\mathcal{G}\cos\left(\frac{\phi_1^+ + \phi_2^+}{2}\right),$

$$\Downarrow$$

$$\psi_-^{(+)} = \lambda\zeta_1^-\cos\left(\frac{\varphi_+^{(+)}}{2}\right), \qquad F_-^{(-)} = \lambda q_2^-\cos\left(\frac{\varphi_+^{(+)}}{2}\right),$$

$$\partial_z\varphi_-^{(-)} = \frac{\lambda}{2}\sin\left(\frac{\varphi_+^{(+)}}{2}\right)\zeta_1^-\psi_+^{(-)} + \lambda\partial_z q_1^-\cos\left(\frac{\varphi_+^{(+)}}{2}\right).$$

- $\bar{D}_-\phi_1^- = -\bar{D}_-\phi_2^- - \frac{8}{\lambda}\mathcal{F}\cos\left(\frac{\phi_1^+ - \phi_2^+}{2}\right),$

$$\Downarrow$$

$$\bar{\psi}_+^{(+)} = -\frac{8}{\lambda}\xi_2^-\cos\left(\frac{\varphi_-^{(+)}}{2}\right), \qquad F_+^{(-)} = -\frac{8}{\lambda}p_2^-\cos\left(\frac{\varphi_-^{(+)}}{2}\right),$$

$$\partial_{\bar{z}}\varphi_+^{(-)} = -\frac{4}{\lambda}\sin\left(\frac{\varphi_-^{(+)}}{2}\right)\xi_2^-\bar{\psi}_-^{(-)} - \frac{8}{\lambda}\partial_{\bar{z}}p_1^-\cos\left(\frac{\varphi_-^{(+)}}{2}\right).$$

Appendix B

Applying the permutability theorem to eqns. (22) and (24) after neglecting the contribution proportional to fermionic superfields, we obtain the following relations

$$\begin{aligned} \gamma_1 s^+_{0,1} c^-_{0,1} + \gamma_2 s^+_{1,3} c^-_{1,3} &= \gamma_2 s^+_{0,2} c^-_{0,2} + \gamma_1 s^+_{2,3} c^-_{2,3}, \\ \gamma_1 s^-_{0,1} c^+_{0,1} + \gamma_2 s^-_{1,3} c^+_{1,3} &= \gamma_2 s^-_{0,2} c^+_{0,2} + \gamma_1 s^-_{2,3} c^+_{2,3}. \end{aligned}$$

Summing and subtracting the above eqns., we find

$$\begin{aligned} &\gamma_1 \left[(s^+_{0,1} c^-_{0,1} \pm s^-_{0,1} c^+_{0,1}) - (s^+_{2,3} c^-_{2,3} \pm s^-_{2,3} c^+_{2,3}) \right] \\ &+\gamma_2 \left[(s^+_{1,3} c^-_{1,3} \pm s^-_{1,3} c^+_{1,3}) - (s^+_{0,2} c^-_{0,2} \pm s^-_{0,2} c^+_{0,2}) \right] = 0. \end{aligned} \tag{35}$$

Using the identity

$$\sin a \cos b \pm \sin b \cos a = \sin\left(a \pm b\right), \tag{36}$$

and eqns. (26) and (27) we can rewrite (35) as

$$\begin{aligned} &\gamma_1 \left\{ \sin\left[\left(\frac{\phi_0^+ + \phi_1^+}{2} \right) \pm \left(\frac{\phi_0^- + \phi_1^-}{2} \right) \right] - \sin\left[\left(\frac{\phi_2^+ + \phi_3^+}{2} \right) \pm \left(\frac{\phi_2^- + \phi_3^-}{2} \right) \right] \right\} \\ &\gamma_2 \left\{ \sin\left[\left(\frac{\phi_1^+ + \phi_3^+}{2} \right) \pm \left(\frac{\phi_1^- + \phi_3^-}{2} \right) \right] - \sin\left[\left(\frac{\phi_0^+ + \phi_2^+}{2} \right) \pm \left(\frac{\phi_0^- + \phi_2^-}{2} \right) \right] \right\} = 0. \end{aligned}$$

Using the fact that

$$\sin a - \sin b = 2 \cos\left(\frac{a+b}{2} \right) \sin\left(\frac{a-b}{2} \right),$$

yields

$$\begin{aligned} 2\cos\left(Y^+ \pm Y^- \right) &\left\{ \gamma_1 \sin\left[(X^+_{1,2} \pm X^-_{1,2}) - (X^+_{3,0} \pm X^-_{3,0}) \right] \right. \\ &\left. +\gamma_2 \sin\left[(X^+_{1,2} \pm X^-_{1,2}) + (X^+_{3,0} \pm X^-_{3,0}) \right] \right\} = 0, \end{aligned}$$

where we have denoted

$$\begin{aligned} Y^\pm &= \frac{\phi_0^\pm + \phi_1^\pm + \phi_2^\pm + \phi_3^\pm}{4}, \\ X^\pm_{j,k} &= \frac{\phi_j^\pm - \phi_k^\pm}{4}. \end{aligned}$$

from where it follows that

$$\begin{aligned} &(\gamma_1 + \gamma_2) \sin\left(X^+_{1,2} \pm X^-_{1,2} \right) \cos\left(X^+_{3,0} \pm X^-_{3,0} \right) = \\ &\quad (\gamma_1 - \gamma_2) \sin\left(X^+_{3,0} \pm X^-_{3,0} \right) \cos\left(X^+_{1,2} \pm X^-_{1,2} \right), \end{aligned}$$

or,

$$\tan\left(X^{+}_{3,0} \pm X^{-}_{3,0}\right) = \left(\frac{\gamma_1+\gamma_2}{\gamma_1-\gamma_2}\right)\tan\left(X^{+}_{1,2} \pm X^{-}_{1,2}\right),$$

and therefore

$$\left(\frac{\phi^{+}_{3}-\phi^{+}_{0}}{4}\right) \pm \left(\frac{\phi^{-}_{3}-\phi^{-}_{0}}{4}\right) = \arctan\left[\delta\tan\left(X^{+}_{1,2} \pm X^{-}_{1,2}\right)\right],$$

where $\delta = \left(\frac{\gamma_1+\gamma_2}{\gamma_1-\gamma_2}\right)$. Adding and subtracting the above expressions we obtain

$$\phi^{\pm}_{3} = \phi^{\pm}_{0} + \Gamma_{\pm},$$

with

$$\Gamma_{\pm} = 2\arctan\left[\delta\tan\left(X^{+}_{1,2} + X^{-}_{1,2}\right)\right] \pm 2\arctan\left[\delta\tan\left(X^{+}_{1,2} - X^{-}_{1,2}\right)\right].$$

Appendix C

Relations (29)and (31) can be written in matrix form,

$$\begin{pmatrix} \mathcal{F}^{(1,3)} \\ \mathcal{F}^{(2,3)} \end{pmatrix} = \frac{1}{Z}\begin{pmatrix} A & -B \\ C & -D \end{pmatrix}\begin{pmatrix} \mathcal{F}^{(0,1)} \\ \mathcal{F}^{(0,2)} \end{pmatrix} \tag{37}$$

where

$$\begin{aligned}
A &= \kappa_2\lambda_2(\bar{c}^{+}_{0,1}c^{-}_{2,3} + \bar{c}^{+}_{2,3}c^{-}_{0,1}), \\
B &= \kappa_2\lambda_1\bar{c}^{+}_{0,2}c^{-}_{2,3} + \kappa_1\lambda_2\bar{c}^{+}_{2,3}c^{-}_{0,2}, \\
C &= \kappa_2\lambda_1\bar{c}^{+}_{1,3}c^{-}_{0,1} + \kappa_1\lambda_2\bar{c}^{+}_{0,1}c^{-}_{1,3}, \\
D &= \kappa_1\lambda_1(\bar{c}^{+}_{0,2}c^{-}_{1,3} + \bar{c}^{+}_{1,3}c^{-}_{0,2}), \\
Z &= \kappa_2\lambda_1\bar{c}^{+}_{1,3}c^{-}_{2,3} - \kappa_1\lambda_2\bar{c}^{+}_{2,3}c^{-}_{1,3}.
\end{aligned} \tag{38}$$

Introduce eqn. (32) into expressions (38). Consider now the following expansions

$$\begin{aligned}
c^{-}_{k,3} &= c_{k,\Gamma_-}\left(1-\frac{\Delta_-^2}{8}\right) - \frac{\Delta_-}{2}s_{k,\Gamma_-}, \\
\bar{c}^{+}_{k,3} &= \bar{c}_{k,\Gamma_-}\left(1-\frac{\Delta_+^2}{8}\right) + \frac{\Delta_+}{2}\bar{s}_{k,\Gamma_+},
\end{aligned}$$

where we have denoted

$$\begin{aligned}
c_{k,\Gamma_-} &= \cos\left(\frac{\phi^{-}_{k}+\phi^{-}_{0}+\Gamma_-}{2}\right) = c^{-}_{k,0}\sigma_+ - s^{-}_{k,0}\rho_-, \\
s_{k,\Gamma_-} &= \sin\left(\frac{\phi^{-}_{k}+\phi^{-}_{0}+\Gamma_-}{2}\right) = s^{-}_{k,0}\sigma_+ + c^{-}_{k,0}\rho_-, \\
\bar{c}_{k,\Gamma_+} &= \cos\left(\frac{\phi^{+}_{k}-\phi^{+}_{0}-\Gamma_+}{2}\right) = \bar{c}^{+}_{k,0}\sigma_- + \bar{s}^{+}_{k,0}\rho_+ \\
\bar{s}_{k,\Gamma_+} &= \sin\left(\frac{\phi^{+}_{k}-\phi^{+}_{0}-\Gamma_+}{2}\right) = \bar{s}^{+}_{k,0}\sigma_- - \bar{c}^{+}_{k,0}\rho_+,
\end{aligned}$$

and

$$\begin{aligned}
\sigma_{\pm} &= \frac{1 \pm \delta^2 \tan\left(\frac{x+y}{4}\right) \tan\left(\frac{x-y}{4}\right)}{\sqrt{1+\delta^2 \tan^2\left(\frac{x+y}{4}\right)}\sqrt{1+\delta^2 \tan^2\left(\frac{x-y}{4}\right)}}, \\
\rho_{\pm} &= \frac{\delta\left[\tan\left(\frac{x+y}{4}\right) \pm \tan\left(\frac{x-y}{4}\right)\right]}{\sqrt{1+\delta^2 \tan^2\left(\frac{x+y}{4}\right)}\sqrt{1+\delta^2 \tan^2\left(\frac{x-y}{4}\right)}}.
\end{aligned}$$

Next, we expand the expressions for A, B, C, D and Z in power series of f obtaining

$$\begin{aligned}
A &= A_0 + \sum_{j,k=1}^{2} A_{j,k} f_{j,k} + \mathcal{O}(f_0), \\
A_0 &= \kappa_2\lambda_2(\bar{c}^{+}_{0,1} c_{2,\Gamma_-} + c^{-}_{0,1}\bar{c}_{2,\Gamma_+}), \\
A_{j,k} &= \frac{1}{2}\kappa_2\lambda_2(c^{-}_{0,1}\bar{s}_{2,\Gamma_+}\Lambda^{+}_{j,k} - \bar{c}^{+}_{0,1} s_{2,\Gamma_-}\Lambda^{-}_{j,k}),
\end{aligned}$$

$$\begin{aligned}
B &= B_0 + \sum_{j,k=1}^{2} B_{j,k} f_{j,k} + \mathcal{O}(f_0), \\
B_0 &= \kappa_2\lambda_1\bar{c}^{+}_{0,2} c_{2,\Gamma_-} + \kappa_1\lambda_2\bar{c}_{2,\Gamma_+} c^{-}_{0,2}, \\
B_{j,k} &= \frac{1}{2}(\kappa_1\lambda_2 c^{-}_{0,2}\bar{s}_{2,\Gamma_+}\Lambda^{+}_{j,k} - \kappa_2\lambda_1\bar{c}^{+}_{0,2} s_{2,\Gamma_-}\Lambda^{-}_{j,k}),
\end{aligned}$$

$$\begin{aligned}
C &= C_0 + \sum_{j,k=1}^{2} C_{j,k} f_{j,k} + \mathcal{O}(f_0), \\
C_0 &= \kappa_1\lambda_2\bar{c}^{+}_{0,1} c_{1,\Gamma_-} + \kappa_2\lambda_1\bar{c}_{1,\Gamma_+} c^{-}_{0,1}, \\
C_{j,k} &= \frac{1}{2}(\kappa_2\lambda_1 c^{-}_{0,1}\bar{s}_{1,\Gamma_+}\Lambda^{+}_{j,k} - \kappa_1\lambda_2\bar{c}^{+}_{0,1} s_{1,\Gamma_-}\Lambda^{-}_{j,k}),
\end{aligned}$$

$$\begin{aligned}
D &= D_0 + \sum_{j,k=1}^{2} D_{j,k} f_{j,k} + \mathcal{O}(f_0), \\
D_0 &= \kappa_1\lambda_1(\bar{c}^{+}_{0,2} c_{1,\Gamma_-} + c^{-}_{0,2}\bar{c}_{1,\Gamma_+}), \\
D_{j,k} &= \frac{1}{2}\kappa_1\lambda_1(c^{-}_{0,2}\bar{s}_{1,\Gamma_+}\Lambda^{+}_{j,k} - \bar{c}^{+}_{0,2} s_{1,\Gamma_-}\Lambda^{-}_{j,k}),
\end{aligned}$$

$$\begin{aligned}
Z &= Z_0 + \sum_{j,k=1}^{2} Z_{j,k} f_{j,k} + \mathcal{O}(f_0), \\
Z_0 &= \kappa_2\lambda_1\bar{c}_{1,\Gamma_+} c_{2,\Gamma_-} - \kappa_1\lambda_2 c_{1,\Gamma_-}\bar{c}_{2,\Gamma_+}, \\
Z_{j,k} &= \frac{1}{2}(\kappa_2\lambda_1 c_{2,\Gamma_-}\bar{s}_{1,\Gamma_+} - \kappa_1\lambda_2 c_{1,\Gamma_-}\bar{s}_{2,\Gamma_+})\Lambda^{+}_{j,k} \\
&- \frac{1}{2}(\kappa_2\lambda_1 s_{2,\Gamma_-}\bar{c}_{1,\Gamma_+} - \kappa_1\lambda_2 s_{1,\Gamma_-}\bar{c}_{2,\Gamma_+})\Lambda^{-}_{j,k},
\end{aligned}$$

where $\mathcal{O}(f_0)$ denotes terms proportional to f_0. It then follows

$$\frac{X}{Z} \;=\; \frac{X_0}{Z_0}\left[1+\sum_{j,k=1}^{2}\left(\frac{X_{j,k}}{X_0}-\frac{Z_{j,k}}{Z_0}\right)f_{j,k}\right]+\mathcal{O}(f_0),$$

where $X=\{A,B,C,D\}$.

Substituting (37), we obtain

$$\mathcal{F}^{(1,3)} \;=\; \frac{A_0}{Z_0}\mathcal{F}^{(0,1)}-\frac{B_0}{Z_0}\mathcal{F}^{(0,2)}+\omega_1^{(1)}\mathcal{F}^{(0,1)}f_{2,1}+\omega_2^{(1)}\mathcal{F}^{(0,1)}f_{2,2}, \tag{39}$$

$$\mathcal{F}^{(2,3)} \;=\; \frac{C_0}{Z_0}\mathcal{F}^{(0,1)}-\frac{D_0}{Z_0}\mathcal{F}^{(0,2)}+\omega_1^{(2)}\mathcal{F}^{(0,1)}f_{2,1}+\omega_2^{(2)}\mathcal{F}^{(0,1)}f_{2,2}, \tag{40}$$

where

$$\begin{aligned}
\omega_1^{(1)} &= \frac{A_0}{Z_0}\left(\frac{A_{2,1}}{A_0}-\frac{Z_{2,1}}{Z_0}\right)+\frac{B_0}{Z_0}\left(\frac{B_{1,1}}{B_0}-\frac{Z_{1,1}}{Z_0}\right),\\
\omega_2^{(1)} &= \frac{A_0}{Z_0}\left(\frac{A_{2,2}}{A_0}-\frac{Z_{2,2}}{Z_0}\right)+\frac{B_0}{Z_0}\left(\frac{B_{1,2}}{B_0}-\frac{Z_{1,2}}{Z_0}\right),\\
\omega_1^{(2)} &= \frac{C_0}{Z_0}\left(\frac{C_{2,1}}{C_0}-\frac{Z_{2,1}}{Z_0}\right)+\frac{D_0}{Z_0}\left(\frac{D_{1,1}}{D_0}-\frac{Z_{1,1}}{Z_0}\right),\\
\omega_2^{(2)} &= \frac{C_0}{Z_0}\left(\frac{C_{2,2}}{C_0}-\frac{Z_{2,2}}{Z_0}\right)+\frac{D_0}{Z_0}\left(\frac{D_{1,2}}{D_0}-\frac{Z_{1,2}}{Z_0}\right).
\end{aligned}$$

From eqns. (28), we get

$$\begin{aligned}
D_+(\phi_3^+-\phi_0^+) &= \frac{8}{\kappa_1}\mathcal{F}^{(0,1)}c_{0,1}^- + \frac{8}{\kappa_2}\mathcal{F}^{(1,3)}c_{1,3}^-,\\
D_+(\phi_1^+-\phi_2^+) &= \frac{8}{\kappa_1}\mathcal{F}^{(0,1)}c_{0,1}^- - \frac{8}{\kappa_2}\mathcal{F}^{(0,2)}c_{0,2}^-.
\end{aligned}$$

Introducing solution (32) in the first eqn. above, we find

$$\begin{aligned}
D_+(\phi_3^+-\phi_0^+) &= D_+(\Gamma_++\Delta_+)=\\
&= \partial_x\Gamma_+D_+(\phi_1^+-\phi_2^+)+D_+\Delta_+=\\
&= \frac{8}{\kappa_1}\mathcal{F}^{(0,1)}c_{0,1}^- + \frac{8}{\kappa_2}\mathcal{F}^{(1,3)}c_{1,3}^-.
\end{aligned}$$

Using eqn. (39) in the above expression, and taking into account that $\mathcal{F}^{(0,1)}$, $\mathcal{F}^{(0,2)}$, $\mathcal{F}^{(0,1)}f_{2,1}$ and $\mathcal{F}^{(0,1)}f_{2,2}$ are independent, we arrive at the following conditions,

$$\begin{aligned}
&\frac{c_{0,1}^-}{\kappa_1}(\partial_x\Gamma_+-1)-\frac{c_{1,\Gamma_-}}{\kappa_2}\frac{A_0}{Z_0}+\frac{gs_{0,2}^-}{4\kappa_2}\Lambda_{1,2}^+ + \frac{gs_{0,1}^-}{4\kappa_1}\Lambda_{1,1}^+=0,\\
&\frac{c_{0,2}^-}{\kappa_2}\partial_x\Gamma_+-\frac{c_{1,\Gamma_-}}{\kappa_2}\frac{B_0}{Z_0}-\frac{gs_{0,1}^-}{4\kappa_1}\Lambda_{2,1}^+ - \frac{gs_{0,2}^-}{4\kappa_2}\Lambda_{2,2}^+=0,\\
&\frac{c_{0,2}^-}{\kappa_2}\partial_x\Lambda_{1,1}^+ + \frac{c_{0,1}^-}{\kappa_1}\partial_x\Lambda_{2,1}^+ + \frac{gs_{0,2}^-}{4\kappa_2}\Lambda_0^+ - \frac{c_{1,\Gamma_-}}{\kappa_2}\omega_1^{(1)} + \frac{s_{1,\Gamma_-}}{2\kappa_2}\left(\frac{A_0}{Z_0}\Lambda_{2,1}^- + \frac{B_0}{Z_0}\Lambda_{1,1}^-\right)=0,\\
&\frac{c_{0,2}^-}{\kappa_2}\partial_x\Lambda_{1,2}^+ + \frac{c_{0,1}^-}{\kappa_1}\partial_x\Lambda_{2,2}^+ - \frac{gs_{0,1}^-}{4\kappa_1}\Lambda_0^+ - \frac{c_{1,\Gamma_-}}{\kappa_2}\omega_2^{(1)} + \frac{s_{1,\Gamma_-}}{2\kappa_2}\left(\frac{A_0}{Z_0}\Lambda_{2,2}^- + \frac{B_0}{Z_0}\Lambda_{1,2}^-\right)=0.
\end{aligned} \tag{41}$$

Moreover, the chirality condition on (32) gives

$$\bar{D}_-(\phi_3^+ - \phi_0^+) = \bar{D}_-(\Gamma_+ + \Delta_+) = 0,$$

from where we obtain the following eqns.

$$\begin{aligned}
&\frac{\bar{c}_{0,1}^+}{\lambda_1}\partial_y\Gamma_+ + \frac{g\bar{s}_{0,1}^+}{4\lambda_1}\Lambda_{1,1}^+ + \frac{g\bar{s}_{0,2}^+}{4\lambda_2}\Lambda_{1,2}^+ = 0,\\
&\frac{\bar{c}_{0,2}^+}{\lambda_2}\partial_y\Gamma_+ - \frac{g\bar{s}_{0,1}^+}{4\lambda_1}\Lambda_{2,1}^+ - \frac{g\bar{s}_{0,2}^+}{4\lambda_2}\Lambda_{2,2}^+ = 0,\\
&\frac{\bar{c}_{0,2}^+}{\lambda_2}\partial_y\Lambda_{1,1}^+ + \frac{\bar{c}_{0,1}^+}{\lambda_1}\partial_y\Lambda_{2,1}^+ + \frac{g\bar{s}_{0,2}^+}{4\lambda_2}\Lambda_0^+ = 0,\\
&\frac{\bar{c}_{0,2}^+}{\lambda_2}\partial_y\Lambda_{1,2}^+ + \frac{\bar{c}_{0,1}^+}{\lambda_1}\partial_y\Lambda_{2,2}^+ - \frac{g\bar{s}_{0,1}^+}{4\lambda_1}\Lambda_0^+ = 0.
\end{aligned} \tag{42}$$

The two sets of eqns. namely, (41) and (42) give the following solutions,

$$\begin{aligned}
\Lambda_{1,1}^+ &= \Lambda_{2,2}^+ = -\frac{8\mu_-}{g\eta_+\eta_-}\cos\left(\frac{x}{2}\right)\sin\left(\frac{y}{2}\right),\\
\Lambda_{1,2}^+ &= \frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\lambda_2}{\lambda_1}\right)\sin\left(\frac{y}{2}\right),\\
\Lambda_{2,1}^+ &= \frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\lambda_1}{\lambda_2}\right)\sin\left(\frac{y}{2}\right),\\
\Lambda_0^+ &= -\frac{32\mu_-}{(g\eta_+\eta_-)^2}\sin\left(\frac{x}{2}\right)\left[\cos\left(\frac{y}{2}\right)(a+\cos x-\cos y) - 2\mu_+\cos\left(\frac{x}{2}\right)\right],
\end{aligned}$$

where

$$\begin{aligned}
\mu_\pm &= \frac{\gamma_1}{\gamma_2} \pm \frac{\gamma_2}{\gamma_1},\\
a &= \frac{1}{2}\left(\frac{\gamma_1^2}{\gamma_2^2} + \frac{\gamma_2^2}{\gamma_1^2}\right) + 3,\\
\eta_\pm &= \mu_+ - 2\cos\left(\frac{x\pm y}{2}\right).
\end{aligned} \tag{43}$$

In order to determine the coefficients Λ^- we make use of

$$\begin{aligned}
\bar{D}_-(\phi_3^- - \phi_0^-) &= \frac{8}{\lambda_1}\mathcal{F}^{(0,1)}\bar{c}_{0,1}^+ - \frac{8}{\lambda_2}\mathcal{F}^{(1,3)}\bar{c}_{1,3}^+,\\
\bar{D}_-(\phi_1^- - \phi_2^-) &= -\frac{8}{\lambda_1}\mathcal{F}^{(0,1)}\bar{c}_{0,1}^+ + \frac{8}{\lambda_2}\mathcal{F}^{(0,2)}\bar{c}_{0,2}^+,
\end{aligned}$$

which are obtained from (30). Introducing (32) in the first of these eqns. we find

$$\begin{aligned}
\bar{D}_-(\phi_3^- - \phi_0^-) &= \bar{D}_-(\Gamma_- + \Delta_-) =\\
&= \partial_y\Gamma_-\bar{D}_-(\phi_1^- - \phi_2^-) + \bar{D}_-\Delta_- =\\
&= \frac{8}{\lambda_1}\mathcal{F}^{(0,1)}\bar{c}_{0,1}^+ - \frac{8}{\lambda_2}\mathcal{F}^{(1,3)}\bar{c}_{1,3}^+.
\end{aligned}$$

Using eqn. (39) in the above expression and taking into account that $\mathcal{F}^{(0,1)}$, $\mathcal{F}^{(0,2)}$, $\mathcal{F}^{(0,1)}f_{2,1}$ and $\mathcal{F}^{(0,1)}f_{2,2}$ are independent, we arrive at the following expressions,

$$\begin{aligned}
&\frac{\bar{c}^{+}_{0,1}}{\lambda_1}(\partial_y\Gamma_- + 1) - \frac{\bar{c}_{1,\Gamma_+}}{\lambda_2}\frac{A_0}{Z_0} + \frac{g\bar{s}^{+}_{0,1}}{4\lambda_1}\Lambda^-_{1,1} + \frac{g\bar{s}^{+}_{0,2}}{4\lambda_2}\Lambda^-_{1,2} = 0,\\
&\frac{\bar{c}^{+}_{0,2}}{\lambda_2}\partial_y\Gamma_- - \frac{\bar{c}_{1,\Gamma_+}}{\lambda_2}\frac{B_0}{Z_0} - \frac{g\bar{s}^{+}_{0,1}}{4\lambda_1}\Lambda^-_{2,1} - \frac{g\bar{s}^{+}_{0,2}}{4\lambda_2}\Lambda^-_{2,2} = 0,\\
&\frac{\bar{c}^{+}_{0,2}}{\lambda_2}\partial_y\Lambda^-_{1,1} + \frac{\bar{c}^{+}_{0,1}}{\lambda_1}\partial_y\Lambda^-_{2,1} + \frac{g\bar{s}^{+}_{0,2}}{4\lambda_2}\Lambda^-_0 - \frac{\bar{c}_{1,\Gamma_+}}{\lambda_2}\omega^{(1)}_1 - \frac{\bar{s}_{1,\Gamma_+}}{2\lambda_2}\left(\frac{A_0}{Z_0}\Lambda^+_{2,1} + \frac{B_0}{Z_0}\Lambda^+_{1,1}\right) = 0,\\
&\frac{\bar{c}^{+}_{0,2}}{\lambda_2}\partial_y\Lambda^-_{1,2} + \frac{\bar{c}^{+}_{0,1}}{\lambda_1}\partial_y\Lambda^-_{2,2} - \frac{g\bar{s}^{+}_{0,1}}{4\lambda_1}\Lambda^-_0 - \frac{\bar{c}_{1,\Gamma_+}}{\lambda_2}\omega^{(1)}_2 - \frac{\bar{s}_{1,\Gamma_+}}{2\lambda_2}\left(\frac{A_0}{Z_0}\Lambda^+_{2,2} + \frac{B_0}{Z_0}\Lambda^+_{1,2}\right) = 0.
\end{aligned} \tag{44}$$

The chiral condition

$$D_+(\phi^-_3 - \phi^-_0) = D_+(\Gamma_- + \Delta_-) = 0,$$

leads us to

$$\begin{aligned}
&\frac{c^-_{0,1}}{\kappa_1}\partial_x\Gamma_- + \frac{gs^-_{0,1}}{4\kappa_1}\Lambda^-_{1,1} + \frac{gs^-_{0,2}}{4\kappa_2}\Lambda^-_{1,2} = 0,\\
&\frac{c^-_{0,2}}{\kappa_2}\partial_x\Gamma_- - \frac{gs^-_{0,1}}{4\kappa_1}\Lambda^-_{2,1} - \frac{gs^-_{0,2}}{4\kappa_2}\Lambda^-_{2,2} = 0,\\
&\frac{c^-_{0,2}}{\kappa_2}\partial_x\Lambda^-_{1,1} + \frac{c^-_{0,1}}{\kappa_1}\partial_x\Lambda^-_{2,1} + \frac{gs^-_{0,2}}{4\kappa_2}\Lambda^-_0 = 0,\\
&\frac{c^-_{0,2}}{\kappa_2}\partial_x\Lambda^-_{1,2} + \frac{c^-_{0,1}}{\kappa_1}\partial_x\Lambda^-_{2,2} - \frac{gs^-_{0,1}}{4\kappa_1}\Lambda^-_0 = 0.
\end{aligned} \tag{45}$$

Solving (44) and (45) for Λ^-, we find

$$\begin{aligned}
\Lambda^-_{1,1} &= \Lambda^-_{2,2} = \frac{8\mu_-}{g\eta_+\eta_-}\cos\left(\frac{y}{2}\right)\sin\left(\frac{x}{2}\right),\\
\Lambda^-_{1,2} &= -\frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\kappa_2}{\kappa_1}\right)\sin\left(\frac{x}{2}\right),\\
\Lambda^-_{2,1} &= -\frac{8\mu_-}{g\eta_+\eta_-}\left(\frac{\kappa_1}{\kappa_2}\right)\sin\left(\frac{x}{2}\right),\\
\Lambda^-_0 &= -\frac{32\mu_-}{(g\eta_+\eta_-)^2}\sin\left(\frac{y}{2}\right)\left[\cos\left(\frac{x}{2}\right)(a - \cos x + \cos y) - 2\mu_+\cos\left(\frac{y}{2}\right)\right],
\end{aligned}$$

where $\mu_\pm$, a and $\eta_\pm$ are given in (43).

Bibliography

[1] C. Rogers, in "Soliton Theory: a survey of results", Ed. A. Fordy , Manchester Univ. Press. (1990)

[2] H. Wahlquist and F. Estabrook, *Phys. Rev. Lett.* **31** (1973) 1386

[3] Q.P. Liu and Y.F. Xie *Phys. Lett.* **325A** (2004) 139; Q.P. Liu and Xing-Biao Hu *J. Physics* **A38** (2005) 6371

[4] J. F. Gomes, L. H. Ymai and A. H. Zimerman, *Phys. Lett.* **373A** (2009) 1401, [arXiv:0902.2456 [math-ph]]

[5] T. Inami and Kanno, *Nucl. Phys.* **B359** (1990) 201; T. Kobayashi and T. Uematsu, *Phys. Lett.* **264B** (1991) 107

[6] "N = 2 and N = 4 supersymmetric mKdV and sinh-Gordon hierarchies"
H. Aratyn, J. F. Gomes, L. H. Ymai and A. H. Zimerman
arXiv:hep-th/0409171

[7] H. Aratyn, J. F. Gomes, L. H. Ymai and A. H. Zimerman, *J. Physics* **A41** (2008) 312001, [arXiv:0712.0626 [nlin.SI]]

[8] J. F. Gomes, L. H. Ymai and A. H. Zimerman, JHEP **0803**, 001 (2008), [arXiv:0710.1391 [hep-th]];
J. F. Gomes, L. H. Ymai and A. H. Zimerman, J. Phys. Conf. Ser. **128**, 012004 (2008), [arXiv:0708.2407 [nlin.SI]]

[9] M. Grigoriev and A. Tseytlin, *Int. J. Mod. Phys.* **A23** (2008) 2107, arXiv:0806.2623 [hep-th]

[10] M. Grigoriev and A. Tseytlin, *Nucl. Phys.* **B800** (2008) 450, arXiv:0711.0155 [hep-th]